



Programme Area: Bioenergy

Project: Characterisation of Feedstocks

Title: D6 Final Report (Phase 1) Appendix 8 Part 3

Abstract:

The primary objective of this 2015/16/17 Project was to provide an understanding of UK produced biomass properties, how these vary and what causes this variability.

This document is one of the appendices to the Final Report from the first Phase (2015/16) of the Characterisation of Feedstocks (CofF) project, Deliverable D6. D6 is provided in a number of parts consisting of the main body text plus 13 Appendices, provided in 17 files. These 13 appendices are provided in 12 pdf files plus 46 data files in Microsoft Excel format. The purpose of this report plus its related parts is to report the variability in feedstock properties of UK produced energy biomass, the causes of these variations and the relationship between the feedstock properties and the provenance data collected. Five feedstocks were studied: Miscanthus, willow short rotation coppice (SRC), poplar SRC, poplar grown as short rotation forests (SRF), and spruce SRF, with poplar and Sitka spruce selected to represent broadleaved and coniferous biomass crops respectively. Provenance data include site properties (such as general climate zone and soil chemistry), the conditions at the time of sample collection, and past management of the site and crop with soil samples also collected for analysis. The feedstock samples were analysed in UKAS accredited laboratories.

Context:


The Characterisation of Feedstocks project provides an understanding of UK produced 2nd generation energy biomass properties, how these vary and what causes this variability. In this project, several types of UK-grown biomass, produced under varying conditions, were sampled. The biomass sampled included Miscanthus, Short Rotation Forestry (SRF) and Short Rotation Coppice (SRC) Willow. The samples were tested to an agreed schedule in an accredited laboratory. The results were analysed against the planting, growing, harvesting and storage conditions (i.e. the provenance) to understand what impacts different production and storage methods have on the biomass properties. The main outcome of this project is a better understanding of the key characteristics of UK biomass feedstocks (focusing on second generation) relevant in downstream energy conversion applications, and how these characteristics vary by provenance.

Disclaimer:

The Energy Technologies Institute is making this document available to use under the Energy Technologies Institute Open Licence for Materials. Please refer to the Energy Technologies Institute website for the terms and conditions of this licence. The Information is licensed 'as is' and the Energy Technologies Institute excludes all representations, warranties, obligations and liabilities in relation to the Information to the maximum extent permitted by law. The Energy Technologies Institute is not liable for any errors or omissions in the Information and shall not be liable for any loss, injury or damage of any kind caused by its use. This exclusion of liability includes, but is not limited to, any direct, indirect, special, incidental, consequential, punitive, or exemplary damages in each case such as loss of revenue, data, anticipated profits, and lost business. The Energy Technologies Institute does not guarantee the continued supply of the Information. Notwithstanding any statement to the contrary contained on the face of this document, the Energy Technologies Institute confirms that the authors of the document have consented to its publication by the Energy Technologies Institute.

Appendix 8 Part 3: *Miscanthus* Infield Variation Charts

Moisture Content of *Miscanthus* IFV




Net Calorific Value of *Miscanthus* IFV




Gross Calorific Value (DAF) of *Miscanthus* IFV




Dry Ash content of *Miscanthus* IFV




Dry nitrogen content of *Miscanthus* IFV




Dry chlorine content of *Miscanthus* IFV




Sulphur content of *Miscanthus* IFV




Barium content of *Miscanthus* IFV




Chromium content of *Miscanthus* IFV




Copper content of *Miscanthus* IFV




Molybdenum content of *Miscanthus* IFV




Nickel content of *Miscanthus* IFV




Vanadium content of *Miscanthus* IFV




Zinc content of *Miscanthus* IFV




Arsenic content of *Miscanthus* IFV




Antimony content of *Miscanthus* IFV




Mercury content of *Miscanthus* IFV




Cadmium content of *Miscanthus* IFV




Lead content of *Miscanthus* IFV




Fluorine content of *Miscanthus* IFV




Bromine content of *Miscanthus* IFV




Variation in ash composition of *Miscanthus* IFV




Aluminium content of *Miscanthus* IFV




Calcium content of *Miscanthus* IFV




Iron content of *Miscanthus* IFV




Potassium content of *Miscanthus* IFV




Magnesium content of *Miscanthus* IFV




Manganese content of *Miscanthus* IFV




Sodium content of *Miscanthus* IFV



Phosphorous content of *Miscanthus* IFV



Silicon content of *Miscanthus* IFV



Titanium content of *Miscanthus* IFV

